

23. Which of the following is the solution to the differential equation  $\frac{dy}{dx} = \frac{2xy}{x^2 + 1}$  whose graph contains the point  $(0, 1)$ ?

(A)  $y = e^{x^2}$

(B)  $y = x^2 + 1$

(C)  $y = \ln(x^2 + 1)$

(D)  $y = 1 + \ln(x^2 + 1)$

(E)  $y = \sqrt{1 + 2\ln(x^2 + 1)}$

$$\begin{aligned}u &= x^2 + 1 \\du &= 2x \, dx \\ \int \frac{1}{u} du\end{aligned}$$

$$\frac{1}{y} dy = \frac{2x}{x^2 + 1} dx$$

$$\ln|y| = \ln|x^2 + 1| + C$$

$$\ln 1 = \ln 1 + C \quad C = 0$$

$$\ln|y| = \ln|x^2 + 1|$$

$$y = x^2 + 1$$

24. Sand is deposited into a pile with a circular base. The volume  $V$  of the pile is given by  $V = \frac{r^3}{3}$ , where  $r$  is the radius of the base, in feet. The circumference of the base is increasing at a constant rate of  $5\pi$  feet per hour. When the circumference of the base is  $8\pi$  feet, what is the rate of change of the volume of the pile, in cubic feet per hour?

- (A)  $\frac{8}{\pi}$       (B) 16      (C) 40      (D)  $40\pi$       (E)  $80\pi$

$$\begin{aligned}
 V &= \frac{1}{3} r^3 \\
 \frac{dV}{dt} &= r^2 \frac{dr}{dt} \\
 &= 16 \left( \frac{5}{2} \right) \\
 &= 40
 \end{aligned}$$

$$\begin{aligned}
 C &= 2\pi r \\
 \frac{dC}{dt} &= 2\pi \frac{dr}{dt} \\
 5\pi &= 2\pi \frac{dr}{dt} \\
 \frac{5}{2} &= \frac{dr}{dt}
 \end{aligned}$$

$$\begin{aligned}
 8\pi &= 2\pi r \\
 r &= \frac{8\pi}{2\pi} = 4 \\
 &= \frac{4}{1}
 \end{aligned}$$

25.  $\lim_{h \rightarrow 0} \frac{e^{-1-h} - e^{-1}}{h}$  is

(A) -1

(B)  $-\frac{1}{e}$

(C) 0

(D)  $\frac{1}{e}$

(E) nonexistent

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f'(x)$$

$$\frac{e^{-1} - e^{-1}}{0} \quad \frac{0}{0}$$

$$\frac{-e^{-1-h}}{1} \quad -e^{-1} \quad -\frac{1}{e}$$

26. Let  $f$  be the function given by  $f(x) = x^3 + 5x$ . For what value of  $x$  in the closed interval  $[1, 3]$  does the instantaneous rate of change of  $f$  equal the average rate of change of  $f$  on that interval?

- (A)  $\sqrt{\frac{7}{3}}$  (B)  $\sqrt{\frac{13}{3}}$  (C)  $\sqrt{5}$  (D)  $\sqrt{6}$  (E)  $\sqrt{\frac{19}{3}}$

$$\frac{27+15}{15}$$

$$3x^2 + 5 = \frac{f(3) - f(1)}{3 - 1}$$

$$3x^2 + 5 = \frac{42 - 6}{2}$$

$$3x^2 + 5 = 18$$

$$3x^2 - 13 = 0$$

$$3x^2 = 13$$

$$x^2 = \frac{13}{3}$$

$$x = \pm \sqrt{\frac{13}{3}}$$

27. If  $e^{xy} - y^2 = e - 4$ , then at  $x = \frac{1}{2}$  and  $y = 2$ ,  $\frac{dy}{dx} =$

(A)  $\frac{e}{4}$

(B)  $\frac{e}{2}$

(C)  $\frac{4e}{8-e}$

(D)  $\frac{4e}{4-e}$

(E)  $\frac{8-4e}{e}$

$$e^{xy} \left[ x \frac{dy}{dx} + y \right] - 2y \frac{dy}{dx} = 0$$

$$x e^{xy} \frac{dy}{dx} + y e^{xy} - 2y \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{-y e^{xy}}{x e^{xy} - 2y}$$

$$\frac{-(2)e}{\frac{1}{2}e - 4}$$

$$\frac{-2e}{\frac{1}{2}e - 4}$$

$$\frac{-4e}{e - 8}$$

$$\frac{4e}{8-e}$$

28. Let  $f$  be the function defined by  $f(x) = x^3 + x^2 + x$ . Let  $g(x) = f^{-1}(x)$ , where  $g(3) = 1$ . What is the value of  $g'(3)$ ?

(A)  $\frac{1}{39}$

(B)  $\frac{1}{34}$

(C)  $\frac{1}{6}$

(D)  $\frac{1}{3}$

(E) 39

$$g'(3) = (f^{-1})'(3)$$

if  $(3, 1) \in g$   
 $(1, 3) \in g^{-1}$   
 $(1, 3) \in f$

$$f'(x) = 3x^2 + 2x + 1$$


$$f'(1) = 6$$

$$\frac{1}{6}$$

if  $(c, d)$  on  $f \rightarrow$

$$(f^{-1})'(d) = \frac{1}{f'(c)}$$

$$f'(1)$$



Graph of  $f$

76. The graph of a function  $f$  is shown above. Which of the following limits does not exist?

(A)  $\lim_{x \rightarrow 1^-} f(x)$

0

(B)  $\lim_{x \rightarrow 1} f(x)$

0

(C)  $\lim_{x \rightarrow 3^-} f(x)$

2

(D)  $\lim_{x \rightarrow 3} f(x)$

Does not exist

(E)  $\lim_{x \rightarrow 5} f(x)$

77. Let  $f$  be a function that is continuous on the closed interval  $[1, 3]$  with  $f(1) = 10$  and  $f(3) = 18$ . Which of the following statements must be true?

(A)  $10 \leq f(2) \leq 18$

(B)  $f$  is increasing on the interval  $[1, 3]$ .

(C)  $f(x) = 17$  has at least one solution in the interval  $[1, 3]$ .

(D)  $f'(x) = 8$  has at least one solution in the interval  $(1, 3)$ .

(E)  $\int_1^3 f(x) dx > 20$



78. Let  $R$  be the region bounded by the graphs of  $y = e^x$ ,  $y = e^3$ , and  $x = 0$ . Which of the following gives the volume of the solid formed by revolving  $R$  about the line  $y = -1$ ?


(A)  $\pi \int_0^3 (e^3 - e^x + 1)^2 dx$

(B)  $\pi \int_0^3 (e^3 - e^x - 1)^2 dx$

(C)  $\pi \int_0^3 [(e^3 - e^x)^2 + 1] dx$

(D)  $\pi \int_0^3 [(e^3 - e^x)^2 - 1] dx$

(E)  $\pi \int_0^3 [(e^3 + 1)^2 - (e^x + 1)^2] dx$



$$V = \pi \int_0^3 [(1+e^3)^2 - (1+e^x)^2] dx$$

79. The number of people who have entered a museum on a certain day is modeled by a function  $f(t)$ , where  $t$  is measured in hours since the museum opened that day. The number of people who have left the museum since it opened that same day is modeled by a function  $g(t)$ . If  $f'(t) = 380(1.02^t)$  and  $g'(t) = 240 + 240\sin\left(\frac{\pi(t-4)}{12}\right)$ , at what time  $t$ , for  $1 \leq t \leq 11$ , is the number of people in the museum at a maximum?

(A) 1      (B) 7.888      (C) 9.446      (D) 10.974      (E) 11

$$f(t) - g(t)$$

$$f'(t) - g'(t) = 0$$

$y^1 \quad y^2$

$y^3 = y^1(x) - y^2(x)$  window  $1 \rightarrow 11$   
 look to see where above to below  
 & find that zero

$x$	0	1	2	3
$f(x)$	5	2	3	6
$f'(x)$	-3	1	3	4


80. The derivative of the function  $f$  is continuous on the closed interval  $[0, 4]$ . Values of  $f$  and  $f'$  for selected values of  $x$  are given in the table above. If  $\int_0^4 f'(t) dt = 8$ , then  $f(4) =$

- (A) 0      (B) 3      (C) 5      (D) 10      (E) 13

$$\int_0^4 f'(t) dt = f(4) - f(0)$$

$$f(4) - 5 = 8$$

$$f(4) = 13$$



81. A slope field for a differential equation is shown in the figure above. If  $y = f(x)$  is the particular solution to the differential equation through the point  $(-1, 2)$  and  $h(x) = 3x \cdot f(x)$ , then  $h'(-1) =$

- (A) -6      (B) -2      (C) 0      (D) 1      (E) 12

$$h'(x) = (3x) f'(x) + f(x) (3)$$

$$\begin{aligned} h'(-1) &= (-3) f'(-1) + f(-1) (3) \\ &= (-3) (-2) + (2)(3) \\ &= 12 \end{aligned}$$