

- (b) For $y \leq 11$, find the y -coordinate of each point on the graph where the line tangent to the graph at that point is vertical.

$$1 - 5 \sin y = 0$$
$$y = .201 \text{ or } y = 6.485 \text{ or } y = 9.223.$$

.201
~~2.540~~
6.415
9.223

- (c) Find the average value of the x -coordinates of the points on the graph in the first quadrant between $y = 5$ and $y = 9$.

$$x^2 = -2 + y + 5 \cos y$$

$$x = \sqrt{-2 + y + 5 \cos y}$$

$$\frac{1}{9-5} \int_5^9 \sqrt{-2 + y + 5 \cos y} \, dy = 2.550$$

t (seconds)	0	3	5	8	12
$k(t)$ (feet per second)	0	5	10	20	24


3. Kathleen skates on a straight track. She starts from rest at the starting line at time $t = 0$. For $0 < t \leq 12$ seconds, Kathleen's velocity k , measured in feet per second, is differentiable and increasing. Values of $k(t)$ at various times t are given in the table above.
- (a) Use the data in the table to estimate Kathleen's acceleration at time $t = 4$ seconds. Show the computations that lead to your answer. Indicate units of measure.

$$a(4) = v'(4) = k'(4) \approx \frac{k(5) - k(3)}{5 - 3} = \frac{10 - 5}{2} = \frac{5}{2}$$

$$\therefore \frac{5}{2} \text{ ft/sec}^2$$

t (seconds)	0	3	5	8	12
$k(t)$ (feet per second)	0	5	10	20	24

- (b) Use a right Riemann sum with the four subintervals indicated by the data in the table to approximate $\int_0^{12} k(t) dt$. Indicate units of measure. Is this approximation an overestimate or an underestimate for the value of $\int_0^{12} k(t) dt$? Explain your reasoning.



$$\int_0^{12} k(t) dt = (5)(2) + (10)(3) + (20)(3) + (24)(4) = 191$$

This est. is an overestimate
because $k(t)$ is strictly increasing
on $(0, 12)$

- (c) Nathan skates on the same track, starting 5 feet ahead of Kathleen at time $t = 0$. Nathan's velocity, in feet per second, is given by $n(t) = \frac{150}{t+3} - 50e^{-t}$. Write, but do not evaluate, an expression involving an integral that gives Nathan's distance from the starting line at time $t = 12$ seconds.

$$5 + \int_0^{12} n(t) dt$$

(d) Write an expression for Nathan's acceleration in terms of t .

$$150(t+3)^{-1}$$


$$h(t) = \frac{150}{t+3} - 50e^{-t}$$

$$h'(t) = -150(t+3)^{-2} - 50e^{-t}(-1)$$

$$= -\frac{150}{(t+3)^2} + 50e^{-t}$$

4. Consider the differential equation $\frac{dy}{dx} = \frac{x(y-1)}{4}$.

(a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.



- (b) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(1) = 3$. Write an equation for the line tangent to the graph of f at the point $(1, 3)$ and use it to approximate $f(1.4)$.

$$\left. \frac{dy}{dx} \right|_{(1,3)} = \frac{1}{2}$$

$$y - 3 = \frac{1}{2}(x - 1)$$

$$y = 3 + \frac{1}{2}(x - 1)$$

$$y(1.4) = 3 + \frac{1}{2}(1.4 - 1) = 3.2$$

(c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(1) = 3$.

$$\frac{dy}{dx} = \frac{x(y-1)}{4}$$

$$\frac{1}{y-1} dy = \frac{1}{4} x dx$$

$$\ln|y-1| = \frac{1}{8}x^2 + C$$

$$y-1 = e^{\frac{1}{8}x^2 + C}$$

$$y-1 = Ae^{\frac{1}{8}x^2}$$


$$\text{for } y(1) = 3$$

$$A = \frac{2}{e^{\frac{1}{8}}}$$

$$y = 1 + 2e^{-\frac{1}{8}} e^{\frac{1}{8}x^2}$$

$$y = 1 + 2e^{\frac{1}{8}x^2 - \frac{1}{8}}$$


$$y = 1 + 2e^{\frac{x^2-1}{8}}$$



5. Let R be the region in the first quadrant enclosed by the graphs of $g(x) = \sqrt{x}$ and $h(x) = \frac{x}{3}$, as shown in the figure above.

(a) Find the area of region R .

$$\begin{aligned}
 A &= \int_0^9 \left(\sqrt{x} - \frac{1}{3}x \right) dx = \left[\frac{2}{3}x^{3/2} - \frac{1}{6}x^2 \right]_0^9 \\
 &= \left(18 - \frac{9}{6} \right) - \left(0 \right) \\
 &= \frac{9}{2}
 \end{aligned}$$



- (b) Write, but do not evaluate, an expression involving one or more integrals that gives the volume of the solid generated when R is revolved about the horizontal line $y = 4$.

$$V = \pi \int_0^9 \left[\left(4 - \frac{x}{3}\right)^2 - \left(4 - \sqrt{x}\right)^2 \right] dx$$

- (c) Find the maximum vertical distance between the graph of g and the graph of h between $x = 0$ and $x = 16$. Justify your answer.

$$D(x) = \sqrt{x} - \frac{1}{3}x$$

$$D'(x) = \frac{1}{2\sqrt{x}} - \frac{1}{3} = \frac{3 - 2\sqrt{x}}{6\sqrt{x}}$$

$$D'(x) = 0 \rightarrow 3 - 2\sqrt{x} = 0$$

$$x = \frac{9}{4}$$

$$D(0) = 0$$

$$D(16) = 4 - \frac{16}{3} = -\frac{4}{3} \quad \left(\frac{4}{3}\right)$$

$$D\left(\frac{9}{4}\right) = \frac{3}{4}$$

Since D is cont $[0, 16]$

by EVT max

dist is $\frac{4}{3}$.

6. Let $g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(a) Find $f(26)$.

$$\begin{aligned} f(26) &= \int_0^{26} 4(x+1)^{-2/3} dx = 12 \sqrt[3]{x+1} \Big|_0^{26} \\ &= (36) - (12) \\ &= 24 \end{aligned}$$

$g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(b) Determine the concavity of the graph of $y = f(x)$ for $x > 0$. Justify your answer.

$$f'(x) = g(x)$$
$$f''(x) = g'(x) = -\frac{8}{3}(x+1)^{-5/3}$$

$$= -\frac{8}{3\sqrt[3]{(x+1)^5}}$$

For $x > 0$ $f''(x) < 0 \therefore f$ is
concave down for $x > 0$.

$g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(c) Let h be the function defined by $h(x) = x - f(x)$. Find the minimum value of h on the interval $0 \leq x \leq 26$.

$$h(x) = x - \int_0^x g(t) dt$$

$$h'(x) = 1 - g'(x)$$

$$h'(x) = 0 \rightarrow g'(x) = 1 \rightarrow 4(x+1)^{-2/3} = 1$$

$$\frac{4}{\sqrt[3]{(x+1)^2}} = 1$$

$$4 = \sqrt[3]{(x+1)^2}$$

$$64 = (x+1)^2$$

$$8 = x+1$$

$$x = 7$$

$$h(0) = 0$$

$$h(7) = 7 - \int_0^7 g(x) dx = -5$$

$$h(26) = 26 - \int_0^{26} g(x) dx = 2$$