

- (b) For $y \leq 11$, find the y -coordinate of each point on the graph where the line tangent to the graph at that point is vertical.

$$1 - 5 \sin y = 0 \rightarrow y = .201 \text{ or } y = 6.485 \text{ or } y = 9.223.$$

- (c) Find the average value of the x -coordinates of the points on the graph in the first quadrant between $y = 5$ and $y = 9$.

$$x^2 = -2 + y + 5 \cos y$$

$$x = \sqrt{-2 + y + 5 \cos y}$$

$$\frac{1}{9-5} \int_5^9 \sqrt{-2 + y + 5 \cos y} \, dy = 2.550.$$

t (seconds)	0	3	5	8	12
$k(t)$ (feet per second)	0	5	10	20	24

3. Kathleen skates on a straight track. She starts from rest at the starting line at time $t = 0$. For $0 < t \leq 12$ seconds, Kathleen's velocity k , measured in feet per second, is differentiable and increasing. Values of $k(t)$ at various times t are given in the table above.
- (a) Use the data in the table to estimate Kathleen's acceleration at time $t = 4$ seconds. Show the computations that lead to your answer. Indicate units of measure.

$$a(4) = v'(4) = k'(4) = \frac{k(5) - k(3)}{5 - 3}$$

$$= \frac{10 - 5}{2}$$

$$= \frac{5}{2}$$

$$\therefore \frac{5}{2} \text{ ft/sec}^2$$

t (seconds)	0	3	5	8	12
$k(t)$ (feet per second)	0	5	10	20	24



- (b) Use a right Riemann sum with the four subintervals indicated by the data in the table to approximate $\int_0^{12} k(t) dt$. Indicate units of measure. Is this approximation an overestimate or an underestimate for the value of $\int_0^{12} k(t) dt$? Explain your reasoning.

$$\int_0^{12} k(t) dt \approx (5)(3) + (10)(2) + (20)(3) + (24)(4) = 191$$

$\therefore 191 \text{ feet}$

Our est. is an overest because k is increasing on $(0, 12)$.

- (c) Nathan skates on the same track, starting 5 feet ahead of Kathleen at time $t = 0$. Nathan's velocity, in feet per second, is given by $n(t) = \frac{150}{t+3} - 50e^{-t}$. Write, but do not evaluate, an expression involving an integral that gives Nathan's distance from the starting line at time $t = 12$ seconds.

$$5 + \int_0^{12} n(t) dt$$

(d) Write an expression for Nathan's acceleration in terms of t .


$$n(t) = 150(t+3)^{-1} - 50e^{-t}$$

$$n'(t) = -150(t+3)^{-2} - 50e^{-t}(-1)$$

$$= -\frac{150}{(t+3)^2} + 50e^{-t}$$

4. Consider the differential equation $\frac{dy}{dx} = \frac{x(y-1)}{4}$.

(a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.



- (b) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(1) = 3$. Write an equation for the line tangent to the graph of f at the point $(1, 3)$ and use it to approximate $f(1.4)$.

$$\left. \frac{dy}{dx} \right|_{(1,3)} = \frac{1}{2}$$

$$y - 3 = \frac{1}{2}(x - 1)$$

$$y = 3 + \frac{1}{2}(x - 1)$$

$$f(1.4) \approx y(1.4) = 3 + \left(\frac{1}{2}\right)(.4) = 3.2$$

(c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(1) = 3$.

$$\frac{dy}{dx} = \frac{x(y-1)}{4}$$

$$\frac{1}{y-1} dy = \frac{1}{4} x dx$$

$$\ln|y-1| = \frac{1}{8} x^2 + C$$

$$y-1 = e^{\frac{1}{8} x^2 + C}$$

$$y = 1 + Ae^{\frac{1}{8} x^2}$$

$$3 = 1 + Ae^{\frac{1}{8}}$$


$$2 = Ae^{\frac{1}{8}} \rightarrow A = \frac{2}{e^{\frac{1}{8}}}$$

$$y = 1 + 2e^{-\frac{1}{8}} e^{\frac{1}{8} x^2} = 2e^{-\frac{1}{8}}$$

$$y = 1 + 2e^{\frac{1}{8} x^2 - \frac{1}{8}}$$

$$y = 1 + 2e^{\frac{x^2 - 1}{8}}$$

$$y = 1 + 2e^{\frac{x^2 - 1}{8}}$$



$$\sqrt{x^3}$$

$$\sqrt{9^3}$$


5. Let R be the region in the first quadrant enclosed by the graphs of $g(x) = \sqrt{x}$ and $h(x) = \frac{x}{3}$, as shown in the figure above.

(a) Find the area of region R .

$$A = \int_0^9 \left(\sqrt{x} - \frac{1}{3}x \right) dx = \left[\frac{2}{3}x^{3/2} - \frac{1}{6}x^2 \right]_0^9$$

$$= \left(18 - \frac{9}{6} \right) - (0)$$

$$= \frac{9}{2}$$



- (b) Write, but do not evaluate, an expression involving one or more integrals that gives the volume of the solid generated when R is revolved about the horizontal line $y = 4$.

$$V = \pi \int_0^9 \left[\left(4 - \frac{x}{3} \right)^2 - \left(4 - \sqrt{x} \right)^2 \right] dx$$

- (c) Find the maximum vertical distance between the graph of g and the graph of h between $x = 0$ and $x = 16$. Justify your answer.

$$D(x) = \sqrt{x} - \frac{1}{3}x$$

$$D'(x) = \frac{1}{2\sqrt{x}} - \frac{1}{3} = \frac{3 - 2\sqrt{x}}{6\sqrt{x}}$$

$$D'(x) = 0 \rightarrow \begin{aligned} 3 - 2\sqrt{x} &= 0 \\ 2\sqrt{x} &= 3 \\ x &= \frac{9}{4} \end{aligned}$$

$$D(0) = 0$$

$$D(16) = 4 - \frac{16}{3} = -\frac{4}{3} \left(\frac{9}{3} \right)$$

$$D\left(\frac{9}{4}\right) = \frac{3}{4}$$

Since $D(x)$ is continuous on $(0, 16)$
by EVT max dist
is $\frac{3}{4}$.

6. Let $g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(a) Find $f(26)$.

$$\begin{aligned} f(26) &= \int_0^{26} 4(x+1)^{-2/3} dx = \left[12 \sqrt[3]{x+1} \right]_0^{26} \\ &= (36) - (12) \\ &= 24 \end{aligned}$$

$g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(b) Determine the concavity of the graph of $y = f(x)$ for $x > 0$. Justify your answer.

$$f'(x) = g(x)$$
$$f''(x) = g'(x) = -\frac{8}{3}(x+1)^{-5/3} = -\frac{8}{\sqrt[3]{(x+1)^5}}$$

$$\text{For } x > 0 \rightarrow f''(x) < 0$$

$\therefore f$ is concave down.

$g(x) = 4(x+1)^{-2/3}$ and let f be the function defined by $f(x) = \int_0^x g(t) dt$ for $x \geq 0$.

(c) Let h be the function defined by $h(x) = x - f(x)$. Find the minimum value of h on the interval $0 \leq x \leq 26$.

$$h'(x) = 1 - g(x) = 1 - 4(x+1)^{-2/3}$$

$$h'(x) = 0 \rightarrow 1 = 4(x+1)^{-2/3}$$

$$1 = \frac{4}{\sqrt[3]{(x+1)^2}}$$

$$4 = \sqrt[3]{(x+1)^2}$$

$$64 = (x+1)^2$$


$$x = 7$$

$$h(0) = 0$$

$$h(26) = 2$$

$$h(7) = -5$$

Since h is cont. on $[0, 26]$
the min value of h is -5 .



Possible inflect points at $x=0$, $x=2$ or $x=3$.

g has inflect. pt at $x=0$ because $g'(x) > 0$ on $(-2,0)$
and $g''(x) < 0$ on $(0,2)$