

This exercise is not intended as a
comprehensive
review of the material.

It is a review...but the main purpose is to
practice
our presentation skills.

Limits Test--14 problems--70 total points

7 limit problems

1 HA

1 VA

1 IVT

1 limits from a graph

1 limit proof

1 continuity

1 remove and redfine

1. Evaluate: $\lim_{x \rightarrow 3} \frac{x^2 + 4x - 21}{x - 3}$.

$$\lim_{x \rightarrow 3} \frac{\cancel{(x-3)}(x+7)}{\cancel{x-3}} = 10$$

2. Evaluate: $\lim_{x \rightarrow 0} \frac{\sqrt{4-x} - 2}{x}$.

$$\lim_{x \rightarrow 0} \frac{\sqrt{4-x} - 2}{x} \cdot \frac{\sqrt{4-x} + 2}{\sqrt{4-x} + 2}$$

$$= \lim_{x \rightarrow 0} \frac{\textcircled{4} - \cancel{x} - \textcircled{4}}{\cancel{x}(\sqrt{4-x} + 2)}$$

$$= \frac{-1}{4}$$

3. Evaluate: $\lim_{x \rightarrow 3} \frac{x-6}{x-3}$.

$$\lim_{x \rightarrow 3} \frac{x-6}{x-3} \nexists$$

$$\lim_{x \rightarrow 3^+} \frac{x-6}{x-3} = -\infty \text{ and } \lim_{x \rightarrow 3^-} \frac{x-6}{x-3} = +\infty$$

4. Evaluate: $\lim_{x \rightarrow \infty} \frac{5 - 2x - 7x^3}{2x^3 - 11x}$.

$$\lim_{x \rightarrow \infty} \frac{5 - 2x - 7x^3}{2x^3 - 11x} = -\frac{7}{2}$$

5. Evaluate: $\lim_{x \rightarrow 25} \frac{\sqrt{x} - 5}{x - 25}$.

$$\begin{aligned} & \lim_{x \rightarrow 25} \frac{\sqrt{x} - 5}{x - 25} \cdot \frac{\sqrt{x} + 5}{\sqrt{x} + 5} \\ &= \lim_{x \rightarrow 25} \frac{\cancel{x - 25}}{(\cancel{x - 25})(\sqrt{x} + 5)} \\ &= \frac{1}{10} \end{aligned}$$

6. Evaluate: $\lim_{x \rightarrow 0} \frac{\sin 5x}{\sin 2x}$.

$$\lim_{x \rightarrow 0} \frac{\sin 5x}{\sin 2x} = \frac{5}{2}.$$

OR

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\left(\frac{\sin 5x}{5x} \right) \cdot 5x}{\left(\frac{\sin 2x}{2x} \right) \cdot 2x} \\ = \frac{5}{2}. \end{aligned}$$

7. Evaluate: $\lim_{x \rightarrow -\infty} \frac{\sqrt{5x^2 - 2x}}{3x + 6}$.

$$\lim_{x \rightarrow -\infty} \frac{\sqrt{5x^2 - 2x}}{3x + 6} = -\frac{\sqrt{5}}{3}$$

8. Find the vertical asymptotes (if any) of $f(x) = \frac{x+5}{(x-2)^2}$.

f has a VA at $x=2$ because
 $f(2) \nexists$ and $\lim_{x \rightarrow 2} f(x) = \pm\infty$.

$f \nexists$ at $x=2$

$\lim_{x \rightarrow 2} \frac{x+5}{(x-2)^2} = \pm\infty$

9. Find the horizontal asymptotes (if any) of $g(x) = \frac{5x - 1}{\sqrt{3x^2 + 6}}$.

g has a HA at $y = \frac{5}{\sqrt{3}}$ because $\lim_{x \rightarrow \infty} g(x) = \frac{5}{\sqrt{3}}$.

g has a HA at $y = -\frac{5}{\sqrt{3}}$ because $\lim_{x \rightarrow -\infty} g(x) = -\frac{5}{\sqrt{3}}$.

10. The function $f(x) = \frac{2x^2 + 5x - 7}{x - 1}$ is discontinuous at $x = 1$. Determine if the discontinuity is removable. If removable, redefine f so that it is continuous at $x = 1$.

Removable?

The discon is removable
because $\lim_{x \rightarrow 1} f(x) = 9$.

Redefine f

$$f(x) = \begin{cases} \frac{2x^2 + 5x - 7}{x - 1} & x \neq 1 \\ 9 & x = 1 \end{cases}$$

11. Determine where the following function is discontinuous: $f(x) = \begin{cases} x^2 - 4 & \text{if } x \leq -2 \\ 4 - x^2 & \text{if } -2 < x < 2 \\ 3 + x & \text{if } x \geq 2 \end{cases}$.

Cont test at $x = -2$

$$f(-2) = 0$$

$$\lim_{x \rightarrow -2^+} f(x) = 0$$

$$\lim_{x \rightarrow -2^-} f(x) = 0$$

$$\left. \begin{array}{l} \lim_{x \rightarrow -2^+} f(x) = 0 \\ \lim_{x \rightarrow -2^-} f(x) = 0 \end{array} \right\} \therefore \lim_{x \rightarrow -2} f(x) = 0$$

$\therefore f$ is cont at $x = -2$

$$\text{because } f(-2) = \lim_{x \rightarrow -2} f(x).$$

Cont test at $x = 2$

$$f(2) = 5$$

$$\lim_{x \rightarrow 2^+} f(x) = 5$$

$$\lim_{x \rightarrow 2^-} f(x) = 0$$

$$\left. \begin{array}{l} \lim_{x \rightarrow 2^+} f(x) = 5 \\ \lim_{x \rightarrow 2^-} f(x) = 0 \end{array} \right\} \therefore \lim_{x \rightarrow 2} f(x) \nexists$$

$\therefore f$ is disc at $x = 2$

$$\text{because } \lim_{x \rightarrow 2} f(x) \nexists.$$

12. Given $f(x) = x^3 - 4x^2 + 4x + 3$, use the Intermediate Value Theorem to show that $f(x) = 0$ for some x between $x = -1$ and $x = 2$.

Since $f(-1) = -6 < 0$ and $f(2) = 3 > 0$
then by IVT $f(x) = 0$ for some $x \in (-1, 2)$.

13. Prove: $\lim_{x \rightarrow 2} x^2 + 2x + 1 = 9$

pf: We need to show that $\forall \epsilon > 0 \exists \delta > 0 \ni$
when $|x - 2| < \delta \rightarrow |(x^2 + 2x + 1) - 9| < \epsilon$

$$|x^2 + 2x - 8| < \epsilon$$

$$|x - 2| |x + 4| < \epsilon$$

$$|x - 2| < \frac{\epsilon}{|x + 4|}$$

Consider $[1, 3]$

$$x = 1 \rightarrow \frac{\epsilon}{|x + 4|} = \frac{\epsilon}{5}$$

$$x = 3 \rightarrow \frac{\epsilon}{|x + 4|} = \frac{\epsilon}{7}$$

$$\therefore \text{choose } \delta = \min \left\{ 1, \frac{\epsilon}{7} \right\}$$

